

# CHEST PAIN (CP)

ICD-9 CODE	ICD-9 DESCRIPTION	ICD-10 CODE	ICD-10 DESCRIPTION	DOCUMENTATION CONCEPTS
786.50	<div style="border: 1px dashed red; padding: 5px; text-align: center; color: red;">                     Not a complete list                 </div> <p>Chest pain, unspecified</p>		<div style="background-color: #00b050; color: white; padding: 5px; text-align: center;">                     More specific ICD-10 options include:                 </div> <p>R07.9 Chest pain, unspecified</p> <p>R07.89 Other chest pain</p> <p>R07.2 Precordial pain</p> <p>R07.1 Chest pain on breathing</p> <p>R07.81 Pleurodynia</p> <p>R07.82 Intercostal chest pain</p>	<p>Chest pain is a symptom</p> <p><u>Location / Type:</u></p> <p>Intercostal Musculoskeletal Non-cardiac Precordial, anterior wall, etc. Ischemic</p> <p><u>Link to etiology (due to):</u></p> <p>Chest pain due to (e.g.):</p> <ul style="list-style-type: none"> <li>GERD w/o esophagitis (K21.9)</li> <li>Muscle strain, thorax (S29.11)</li> <li>STEMI (I21.3) / NSTEMI (I21.4)</li> <li>Unstable angina (I20.0)</li> <li>Pneumonia (J18.9)</li> <li>Neoplasm (G89.3), etc.</li> </ul> <p><u>Acuity:</u></p> <p>Acute Chronic Acute on chronic</p>

# CHEST PAIN (CP)

ICD-9 DESCRIPTION	ICD-10 DESCRIPTION
<b>INAPPROPRIATE DOCUMENTATION EXAMPLE</b>	<b>APPROPRIATE DOCUMENTATION EXAMPLE</b>
<p><b>Diagnosis:</b> 786.50 Chest pain, unspecified</p> <p><b>HPI:</b> Patient presented with complaint of chest pain.  <b>Associated symptoms:</b> Indigestion and burning in throat.  <b>Pertinent negatives:</b> No edema, no radiation, no PMH CAD.  <b>Compliance with medication:</b> On no meds.  <b>Compliance with lifestyle:</b> Regular diet.  <b>Social History:</b> Consumes 1-2 cans of beer in the evening.</p> <p><b>Assessment/Plan:</b> Non-cardiac chest pain. EKG WNL. Start Prevacid 60 mg/daily.</p> <div data-bbox="142 896 957 972" style="background-color: red; color: white; padding: 5px; text-align: center;">                     Concept(s) requirement not met: no type, no location, no underlying cause                 </div>	<p><b>Diagnosis:</b> K21.9 GERD without esophagitis</p> <p><b>HPI:</b> Patient presented with complaint of anterior, midsternal chest pain.  <b>Associated symptoms:</b> Pain is intermittent, often postprandial, and associated with burning in throat, especially in reclining position.  <b>Pertinent negatives:</b> No edema, no radiation, no PMH CAD.  <b>Compliance with medication:</b> On no meds.  <b>Compliance with lifestyle:</b> Regular diet.  <b>Social History:</b> Consumes 1-2 cans of beer in the evening.</p> <p><b>Assessment/Plan:</b> Non-cardiac chest pain due to GERD. EKG WNL, no ischemia. Start Prevacid 60 mg/daily. Diet: Low fat diet. Avoid citrus and acidic foods. Consider GI referral for EGD if symptoms persist.</p> <div data-bbox="1089 896 1976 972" style="background-color: #00b050; color: white; padding: 5px; text-align: center;">                     Concept(s) requirement met: type, location, underlying etiology, plan of care                 </div>